



Sedimentation of helminth eggs in low quality water to be used for irrigation in agriculture

Sengupta, Mita Eva; Thamsborg, Stig Milan; Andersen, Thorbjørn Joest; Olsen, Annette; Dalsgaard, Anders

Publication date:
2010

Document version
Early version, also known as pre-print

Citation for published version (APA):
Sengupta, M. E., Thamsborg, S. M., Andersen, T. J., Olsen, A., & Dalsgaard, A. (2010). *Sedimentation of helminth eggs in low quality water to be used for irrigation in agriculture*. Poster session presented at ICOPA XII. The XIIth International Congress of Parasitology, Melbourne, Australia.



DEPARTMENT OF VETERINARY DISEASE BIOLOGY
FACULTY OF LIFE SCIENCES
UNIVERSITY OF COPENHAGEN



Sedimentation of helminth eggs in low quality water to be used for irrigation in agriculture

Mita E. Sengupta^{1*}, Stig M. Thamsborg¹, Thorbjørn Joest Andersen², Annette Olsen¹ & Anders Dalsgaard¹



¹ Department of Veterinary Disease Biology, Faculty of Life Sciences, University of Copenhagen, Dyrlægevej 100, DK-1870 Frederiksberg C, Denmark. *Email: msen@life.ku.dk
² Department of Geography & Geology, University of Copenhagen, Øster Voldgade 10, DK- 1350 Copenhagen K, Denmark.

INTRODUCTION

- Irrigation with low quality water in agriculture is increasing worldwide due to water shortage. Low quality water often contains pathogens like eggs of helminth parasites and irrigation with such water may therefore negatively affect food safety, and health of humans and animals.
- Sedimentation of particles, including eggs, is expected to follow Stoke's law, which implies that sedimentation velocity depends on particle size, differences in density between particles and water, and the viscosity of water.
- The present study aimed to experimentally determine the sedimentation velocity of helminth eggs in tap and waste water.



METHOD




- Eggs of *Oesophagostomum* sp., *Ascaris suum* and *Trichuris suis* were recovered from pig faeces by sieving and flotation.
- 'Owen tubes' were filled with 2L of either tap water or inlet water from a sewage plant (dry matter content of 1860 mg/L), and added 500-600 eggs of each helminth.
- 10 sub-samples (200ml) were collected at varying time intervals (4 min to 22 hrs) from the bottom of the Owen tube and the number of eggs was counted.
- Settling velocity was also determined by means of 'Laser In-Situ Scattering and Transmissometry' (LISST).

$$\text{Stoke's Law: } V_s = \frac{g}{18} \times \frac{d^2}{\eta} \times (\rho_p - \rho_l) \times \eta^{-1}$$

g is gravitational acceleration (9.81 m s⁻²)
 d is particle diameter (m)
 ρ_p is specific density of the particles
 ρ_l is specific density of the liquid
 η is dynamic viscosity of the liquid.

RESULTS



- According to size and density, *Trichuris* eggs were expected to sediment at the fastest rate followed by *Ascaris* and *Oesophagostomum* spp. This was confirmed by the Owen-tube method in tap water but not by LISST.
- The settling velocity for *Ascaris* was faster in waste water as compared to tap water. Most likely because eggs are trapped in large sized flocculated particles which have a faster settling rate.
- In contrast, eggs of *Trichuris* and *Oesophagostomum* spp. seemed to be unaltered/ slowed down in waste water.
- In general, the sedimentation rate is slow for all 3 egg types and if there is a water flow the eggs will not sediment, but be transported along with the water.

CONCLUSION



- It was possible to determine the settling velocity of eggs of 3 different helminths in tap and waste water. Sedimentation is slow compared to inorganic particles.
- Particles in waste water influence the settling velocity of eggs but not in a uniform way.
- The results will form basis for improved sanitation of low quality irrigation water.

ACKNOWLEDGEMENT

The study was supported by the "Safe and High Quality Food Production using Low Quality Waters and Improved Irrigation Systems and Management" project (SAFIR, EU, FOOD-CT-2005023168) funded by the European Commission